

**BELLSOUTH**  
**ORIGINAL**

Kathleen B. Levitz  
Vice President-Federal Regulatory

EX PARTE OR LATE FILED

Suite 900  
1133-21st Street, N.W.  
Washington, D.C. 20036-3351  
202 463-4113  
Fax: 202 463-4198  
Internet: levitz.kathleen@bsc.bls.com

October 5, 1998

**RECEIVED**

**OCT - 5 1998**

FEDERAL COMMUNICATIONS COMMISSION  
OFFICE OF THE SECRETARY

Ms. Magalie Roman Salas  
Secretary  
Federal Communications Commission  
1919 M Street, NW, Room 222  
Washington, D.C. 20554


Re: Written Ex Parte in CC Docket No. 98-56 and  
CC Docket No. 98-121

Dear Ms. Salas:

This is to inform you that BellSouth Corporation has submitted today a written ex parte to the staff of the Common Carrier Bureau's Policy and Program Planning Division. That ex parte presents information relating to methods of estimating variance using replicates, a topic of discussion during a meeting on September 16, 1998, attended by representatives of BellSouth, statisticians from Ernst & Young, and Bureau staff. This information has been submitted in response to a request from the staff of the Common Carrier Bureau.

Pursuant to Section 1.1206(a)(1) of the Commission's rules, we are filing two copies of this notice and that written ex parte presentation. Please associate this notification with the record of CC Docket No. 98-121.

Sincerely,



Kathleen B. Levitz

Attachment

cc: Carol Matthey

**BELLSOUTH**

**Kathleen B. Levitz**  
Vice President-Federal Regulatory

Suite 900  
1133-21st Street, N.W.  
Washington, D.C. 20036-3351  
202 463-4113  
Fax: 202 463-4198  
Internet: levitz.kathleen@bsc.bls.com

October 5, 1998

**RECEIVED**

**OCT - 5 1998**

**FEDERAL COMMUNICATIONS COMMISSION  
OFFICE OF THE SECRETARY**

**Ms. Carol Matthey, Chief  
Policy and Program Planning Division  
Common Carrier Bureau  
Federal Communications Commission  
1919 M Street, N.W.  
Washington, D.C. 20554**

Written Ex Parte in CC Docket No. 98-56 and  
CC Docket No. 98-121

Dear Ms. Matthey:

On September 16, 1998, William Stacy, Ray Lee, Craig Duncan, Jerry Moore, Robert Blau and I, all of BellSouth, participated in a videoconference with your staff. Also participating in that conference were Dr. Fritz Scheuren, Dr. Susan Kinkins and Scott Abbott of Ernst & Young. During that videoconference, there was a discussion of variance estimation using replicates. A staff member, Daniel Shiman asked for references discussing this methodology. In response to Mr. Shiman's request, attached is a paper by Hinkins, Oh and Scheuren entitled Replicate Variance Estimates - Reducing Bias by Using Overlapping Replicates that the authors presented at the 1997 Joint Statistical Meetings. I am also attaching a copy of a letter from Dr. Scheuren that offers another source for additional information about this methodology.

If after reviewing this attachment your staff concludes that it needs additional information, please call me at (202) 463-4113.

In compliance with Section 1.1206(a)(1) of the Commission's rules, we have today filed with the Secretary of the Commission two copies of this written ex parte presentation for both of the dockets identified above and request that it be associated with the record of both dockets.

Sincerely,



Kathleen B. Levitz

Attachment

cc: Jake Jennings

Andrea Kearney  
Daniel Shiman

Florence Setzer

September 25, 1998


Kathy Levitz  
BellSouth  
1133 21<sup>st</sup> Street, NW  
Suite 900  
Washington, DC 20036-3351

Dear Kathy,

At last week's meeting with the FCC, there was a discussion of variance estimation using replicates and Daniel Shiman was interested in references for this methodology.

Enclosed is a paper that we presented at the 1997 Joint Statistical Meetings; it includes a brief description of the general methodology, as well as our proposed improvement. For an in-depth discussion (and examples) of variance estimation based on the concept of replication, we would recommend Chapter 2 in Kirk Wolter's book, which is referenced in this paper.

Sincerely,



Fritz Scheuren, Ph.D.  
Associate National Technical Director  
of Statistical Sampling

Encl.

cc: Ray Lee

## REPLICATE VARIANCE ESTIMATES - REDUCING BIAS BY USING OVERLAPPING REPLICATES

Susan Hinkins and H. Lock Oh, IRS, and Fritz Scheuren, Ernst & Young  
Susan Hinkins, 1122 South 5th Ave., Bozeman, MT 59715

**Key Words:** Repeated Samples; Permanent Random Numbers

### 1. BACKGROUND

A replicate variance estimator can be useful when the form of the estimator is complex or when the sampling distribution is complex. In the example that motivated this work, the estimators are simple totals or means, but the sampling distribution is unwieldy as it involves the probability of being in different strata over time.

The original problem of interest grew out of the use of a permanent random number (PRN) for sample selection in the Statistics of Income (SOI) samples, in particular, the annual sample of corporate tax returns. This is a stratified probability sample designed, in part, to provide cross-sectional estimates of income and tax items for a particular year. Since it is an annual sample, estimates of year-to-year change are also of interest, and for users within the Treasury Department, the primary interest is in modeling economic and tax dynamics over time using the microdata. By using a permanent random number in the sample selection, the year-to-year sample overlap is increased while maintaining the simplicity and validity of the cross-sectional estimation.

Because of the sample overlap, the precision of estimates of year-to-year change may be greatly improved. For variables with high year-to-year correlation, the standard error may be reduced by as much as one half, compared to independent samples. Calculating estimates of variance is more difficult, however, since the probability of a unit being in both samples depends on its sampling stratum each year, and this can change.

Hinkins, Moriarity, and Scheuren

(1996) describe the SOI corporate sample, the selection using a PRN, and the resulting year-to-year overlap. A method is given for defining replicates, using a PRN, so that a unit stays in the same replicate over time. In this way, replicate variance estimation can be used for estimating the variance of estimates of year-to-year change.

One difficulty with the replicate procedure, in general, is that it does not account for the finite population correction (fpc). In the simple random sample case, the replicate variance estimate is an unbiased estimate of the variance of the mean or total only if the fpc can be ignored. The SOI corporate sample has sampling rates as large as .5, in which case the fpc cannot be dismissed.

In this paper we describe a general modification to the usual replicate variance estimator to adjust for the finite population correction. Section 2 gives a brief description of the replicate variance estimator in general and describes the proposed adjustment to the replicate methodology. Section 3 discusses the case where one wants a variance estimate for the original estimator rather than the replicate estimator. An example based on the SOI corporate sample is given. Section 4 briefly summarizes the results and describes future work.

### 2. OVERLAPPING REPLICATE VARIANCE ESTIMATORS

Replicate variance estimators are useful in many cases where the variance calculation is complex, as they only require calculation of the point estimate (mean, ratio, total, etc). Suppose the sample is of size  $n$  where  $n = m \cdot G$ . The dependent random groups variance estimator (eg. Wolter, 1985)

Figure 1. Overlapping Replicates, k-1

A. With t=1	Group	Units																			
	1	x	x	x	x																x
	2	x				x	x	x	x												
	3					x				x	x	x	x								
	4									x				x	x	x	x				
	5																x		x	x	x

B. With t=3	Group	Units																			
	1	x	x	x	x					x							x				x
	2	x				x	x	x	x								x				x
	3	x				x				x	x	x	x								x
	4	x				x				x				x	x	x	x				
	5					x				x							x		x	x	x

is calculated by using a random mechanism to divide the sample into G groups, each of size m. The estimator of interest, say  $\hat{X}$ , is calculated in each group,  $\hat{X}_g$ . The replicate estimator and variance estimator are

$$\hat{X}_r = \frac{1}{G} \sum_{g=1}^G \hat{X}_g$$

$$V_1 = \hat{\text{var}}(\hat{X}_r) = \frac{1}{G(G-1)} \sum_{g=1}^G (\hat{X}_g - \hat{X}_r)^2$$

For estimators of means and totals, the replicate estimator is equal to the original estimator, and with simple random sampling with finite population correction (fpc), we have

$$\hat{X}_r = \hat{X} \quad \text{and} \quad E(V_1) = \frac{\text{var}(\hat{X})}{(fpc)}$$

The replicate variance estimate is a conservative estimate, overestimating the variance, and is approximately unbiased only when the fpc is close to one.

For a stratified sample, if the fpc's were equal for every stratum, one could simply correct the replicate variance estimator. Having nonconstant fpc's across strata would be typical in highly skewed populations and is true in the SOI sample; hence a simple adjustment is unavailable to us.

**Overlapping Replicates.** -- For most situations, there is a reasonably straightforward way to adjust the definition of the replicates in order to get an approximately unbiased estimate of variance. Note that in the case  $n = m \cdot G$ , the expected value of the variance estimator  $V_1$  can be written (e.g., Wolter 1985) as

$$E(V_1) = \text{var}(\hat{X}_r) = \frac{1}{G(G-1)} \sum_{g \neq h} \text{Cov}(\hat{X}_g, \hat{X}_h)$$

If there is no intervention, because of the fpc's, the covariance terms between the estimators from different random groups are all equal and negative, and  $V_1$  will, as a result, be positively biased. What if one could alter the covariance between replicate estimates, so that the total covariance term was approximately zero? Then, the replicate variance estimator  $V_1$  would be nearly unbiased.

Assume that the original sample has been divided into G dependent random groups each of size m. And assume that the groups are randomly ordered and that the units within groups are randomly ordered. Then we can denote the sample and the random groups as n units, 1, 2, ..., n, where the first group consists of units 1 through m, the second group consists of units m+1, m+2, ..., 2m, etc. Figure 1 shows an example with  $n = 20$  and  $G = 5$ ; each

group then has  $n-4$  members.

In general, we want to form  $G$  new groups by randomly selecting  $k$  units in the original group 1 to overlap with the next consecutive  $t$  groups. Then we select  $k$  units from the original group 2 to overlap with the next consecutive  $t$  groups. Only two values of  $t$  need to be considered:  $t=1$  and  $t=3$ .

Figure 1 shows an example for both  $t=1$  and  $t=3$ , using  $k=1$ . In each case there are still 5 replicates, but now each replicate contains  $n + t \cdot k$  units.

With overlapping replicates ( $k > 0$ ), the replicate estimate of the total,  $\hat{X}_r$ , no longer equals the original sample estimate of the total,  $\hat{X}$ . But conditional on the sample achieved,

$$E(\hat{X}_r | \text{sample}) = \hat{X}.$$

However, the replicate variance estimate,  $V_1$ , is now an estimator of the replicate estimate  $\hat{X}_r$ , rather than the original estimate  $\hat{X}$ .

In the case of  $t=3$ , the restriction  $G \geq 5$  and  $m \geq 3$  is needed, which is not an unreasonable requirement for using replicate estimates in general. Then for  $t=1$  or  $t=3$ ,

$$E(V_1) = \text{Var}(\hat{X}_r) = h_t(k) \cdot N \cdot S^2$$

where

$$h_t(k) = \frac{t(t+1)kN}{(G-1)(m+tk)^2} - 1.$$

By solving for the value of  $k$  which makes  $h_t(k) = 0$ , an unbiased estimate  $V_1$  can be constructed.

In the case  $t=1$ , if the sampling rate and the number of replicates,  $G$ , satisfy

$$\frac{n}{N} < \frac{1}{2} + \frac{1}{2(G-1)}$$

then there is a solution

$$k_1 = \frac{N - (n-m) - \sqrt{N(N-2(n-m))}}{G-1}$$

which satisfies  $0 < k \leq m$ . Therefore,

for sampling rates no larger than .5, there is a solution for any value of  $G$ .

Most sampling designs probably fall into this category, i.e. with sampling rates all less than or equal to 0.5. If there are strata that are selected with probability 1.0, then the usual solution is to include the entire certainty stratum in each replicate, as discussed later. For cases with sampling rates between .5 and 1.0, we can use  $t=3$ , and the solution

$$k_1 = \frac{2N - (n-m) - 2\sqrt{N(N-(n-m))}}{3(G-1)}$$

satisfies  $0 < k < n/3$  for all sampling rates.

Choosing  $t=1$  vs  $t=3$  -- Since using  $t=3$  gives a solution for all sampling rates, why bother with the case  $t=1$ ? One reason is that the case  $t=1$  is easier to construct. The second reason is that the case  $t=1$  is more likely to result in a reduction in bias for smaller sampling rates.

At the exact solution  $k$ , we would have an unbiased estimate. However, we get only an approximately unbiased estimate,  $V_1$ , because  $k$  must be rounded to an integer value. In order to assure a conservative estimate of variance, one should always round down. That is, in both cases  $t=1$  and  $t=3$ , one can show that rounding down will result in a negative value of  $h_t(k)$ , whereas rounding up will result in a positive value. So rounding down will result in a (hopefully small) over-estimate of the variance.

Therefore, if the exact solution  $k$  is less than 1, we round down to 0 and we do not reduce the bias. Conditions under which there will be a useful solution can be described in terms of the initial sampling rate,  $f$ , the population size  $N$ , and the number of replicates,  $G$ . Namely, if

$$f \geq G \cdot \left( \sqrt{\frac{f \cdot (f+1)}{N \cdot (G-1)}} - \frac{t}{N} \right)$$

than or equal to 1. One would hope that the value of the right hand side of the inequality would be relatively small. Holding  $N$  and  $G$  fixed, the value of the right hand side is smaller for  $t=1$  than for  $t=3$ . Therefore, for sampling rates of .5 or less, the method of overlapping replicates using  $t=1$  will reduce the bias of the variance estimate for smaller sampling rates compared to overlapping with  $t=3$ .

This is not such an important consideration for large populations. For example, with  $N=100,000$  and  $G=25$ , one can get a bias reduction using  $t=3$  for any design with sampling rate greater than .055. Using  $t=1$ , one can get a bias reduction for designs with sampling rates down to .023. But at such small sampling rates, the bias of the usual replicate variance estimate is very small anyway. However, with smaller populations, the difference can be noticeable. Take for example  $N=10,000$  and  $G=25$ . Using  $t=3$ , one gets a reduction in the bias only for sampling rates larger than .17. Using  $t=1$ , one can reduce the bias for sampling rates as low as .07. If the sampling fraction is .1, using the configuration with  $t=3$  will not result in a bias reduction, but using  $t=1$  will.

In general, the cases where this method does not reduce the bias appear to coincide with examples where the replicate estimate may not be useful in general, namely small sample sizes. When the population size is small, one cannot have both a small sampling rate and a large number of replicates. This does not seem unreasonable; one cannot expect to use the replicate method if the sample is very small.

### 3. REPLICATE VS ORIGINAL ESTIMATOR

For estimation of means or totals, the usual random groups replicate estimator, with no overlap, is the same as the original estimator. In this case,  $V_1$  is an estimator of the variance of the original estimate,  $\bar{X}$ .

When overlapping replicates are used, the replicate estimator,  $\hat{X}_r$ , is no longer equal to the original estimator,  $\bar{X}$ . And the variance of the replicate estimator will be larger than the variance of the original estimator.

This is most immediately noticeable with certainty strata. The variance of the original estimate is zero. By including the entire certainty stratum in each replicate, this property is preserved and we have

$$\hat{X}_r = \bar{X} \quad \text{Var}(\hat{X}_r) = 0 \quad \text{and} \quad V_1 = 0.$$

Note that one could also divide the certainty strata into  $G$  random groups and use the general solution with  $t=3$  to find a value of  $k < m/3$  that results in an unbiased estimator  $\hat{X}_r$  and an approximately unbiased variance estimator,  $V_1$  for  $\hat{X}_r$ . But this is not the best solution for certainty strata, in the sense that

$$\hat{X}_r \neq \bar{X} \quad \text{and} \quad \text{Var}(\hat{X}_r) > \text{Var}(\bar{X}) = 0.$$

In other cases as well, one may want a replicate variance estimate that is an unbiased estimate of the variance of the original sample estimate. This can be done using the fact that

$$\text{Var}(\hat{X}_r) = \text{Var}(\bar{X}) + E(\text{Var}(\hat{X}_r | \text{sample})).$$

In the case  $t=1$ , we find, for totals,

$$E(V_1) = \text{Var}(\bar{X}) - N \left( h_1(k) - \frac{k(m-k)N}{n(m+k)^2} \right) S^2$$

where  $h_1(k)$  was defined in Section 2. By solving for the value of  $k$ , say  $k_2$ , that makes the coefficient on  $S^2$  equal to zero, we have an unbiased estimate. In order for  $0 < k_2 \leq m$ , the same condition as before is required, and the solution in terms of the proportion of overlap is

$$k_2 = \frac{1}{m} \left[ \bar{X} - \frac{(G+1)}{2(G-1)} \left( 1 - \sqrt{1 - \frac{8f(G-1)}{(G+1)^2}} \right) \right]$$

**Table 1. Example of Overlapping Replicates for Stratified Design, G=25, t=1**  
**Relative increase in Var**

Stratum	$N_h$	$n_h$	$m_h$	$k_1$	$k_2$	Using $k_1$	With $k_2$
1	140,000	7,050	282	7	13	.024	.042
2	50,000	2,950	118	3	6	.025	.046
3	28,000	2,950	118	6	12	.049	.084
4	20,000	5,950	238	49	92	.160	.176
5	10,000	5,000	200	133	184	.161	.040

which lies between 0 and 1.

There are several choices here. For a given value of t, either t=1 or t=3, there are three replicate estimators of interest, namely those associated with k=0 (no overlap) or k=k<sub>1</sub> or k=k<sub>2</sub>. With each replicate estimator there is an associated replicate variance estimator, V<sub>i</sub>(k).

Using k=0, the replicate estimator is the same as the original estimator. But the associated variance estimate V<sub>i</sub>(0) can be exceedingly conservative when the sampling rates are not small.

Using k=k<sub>1</sub>, V<sub>i</sub> is an approximately unbiased estimate of the variance of the replicate estimator. It is a conservative estimate of the variance of the original estimator. That is, using exact values of k, we would have

$$E(V_i(k_1)) = \text{Var}(\hat{X}_i(k_1)) \geq \text{Var}(\hat{X}_i)$$

Using k=k<sub>2</sub>, V<sub>i</sub> is an approximately unbiased estimate of the variance of the original estimator. But V<sub>i</sub> will under-estimate the variance of its associated replicate estimator:

$$E(V_i(k_2)) = \text{Var}(\hat{X}_i) \leq \text{Var}(\hat{X}_i(k_2))$$

For best results, one needs to decide on the estimator of interest before determining the amount of overlap in the replicates, or else provide more than one definition of replicates. For a general purpose data base, a reasonable compromise might be

to use the construction with an overlap of k<sub>1</sub> units. Then V<sub>i</sub> is an unbiased estimate of the replicate estimate of the total. And, as we will see in the next example, even though it is a biased (but conservative) estimate of the variance of the original estimator, it can be much better than the usual replicate variance estimator.

An Example from SOI -- Take as an example a simplified version of some of the non-certainty SOI strata for the regular corporations, as shown in Table 1. The second and third columns give the population and sample sizes respectively. Using G=25 replicates, the fourth column shows the resulting original group size, m.

Since the largest sampling rate is .5, we can use the configuration t=1 for all strata. Column 5 gives the value of k<sub>1</sub>, the number to overlap in order to get an approximately unbiased variance estimate of the replicate estimator. Column 6 shows the value of k<sub>2</sub>, the overlap needed in order to use V<sub>i</sub> as an (approximately) unbiased estimate of the original stratified estimate of the total (or mean).

The last two columns show the relative increase in variance, by strata, if we use the replicate estimate of the total, compared to the original weighted stratum estimate. For example, in stratum 3 using overlapping replicates with k=6, the variance of the replicate estimate,  $\hat{X}_i$ ,



is approximately 5% larger than the variance of the usual estimate  $\bar{X}$ .

For  $t=1$ , the maximum increase in variance occurs at  $k = m/3$ . So if  $k_1 < k_2 < m/3$ , using  $k_2$  results in a larger variance of  $\bar{X}$ , than using  $k_1$ . But if, as in stratum 5,  $m/3 < k_1 < k_2$ , then the variance of the replicate estimator using  $k_2$  is smaller than the variance of the replicate estimator using  $k_1$ .

Suppose we are interested in using  $V_1$  as an estimator for the variance of the original stratified estimate of the total. We can calculate the relative bias,  $B$ , of the estimator  $V_1(k)$ , for each value of  $k$ :

$$B(V_1(k)) = \text{Var}(\hat{X}) / (1 + B(k))$$

where  $B$  depends on population and sample sizes, the number of replicates,  $G$ , as well as the size of the overlap,  $k$ . Table 2 shows the relative bias, by stratum. Note that even though  $k_1$  is not optimal, it gives considerably better estimates of the variance than the usual replicate variance estimate ( $k=0$ ), especially when the fpc is not close to one. And it should give approximately unbiased estimates of the replicate estimate of the total.

Table 2. Relative Bias in  $V_1$ , for Estimating  $\bar{X}$ .

Stratum	$k=0$	$k=k_1$	$k=k_2$
1	.053	.025	.003
2	.063	.034	.007
3	.120	.059	.007
4	.420	.164	.003
5	1.000	.161	.0001

The relative bias using  $k=k_2$  should be zero. It is only approximately zero because  $k$  is rounded to an integer. The bias in stratum 5 is so much smaller than the others because in this case the exact value of  $k$  is 184.03, compared to stratum 3, where the exact value of  $k$  is 12.8.

Suppose we decide to define the replicates using  $k_2$ . In each stratum,

we randomly divide the sample units into 25 groups and randomly order the 25 groups. In stratum 1, in each group of 282, we randomly select 13 units and include these in the "next" group as well, etc. Therefore each replicate in stratum 1 will have 295 units; each replicate in stratum 2 will have 124 units, etc. In this way, 25 replicates, each of size 2,021 are formed.

#### 4. CONCLUSIONS AND FUTURE WORK

The results shown here imply that when the fpc factor cannot be ignored, we could improve considerably over the usual dependent group estimates of variance by using overlapping replicates. And this technique is programmable. The results shown here are exact for the relatively unrealistic case where  $n=m+G$ . In practice, we will have some slight variation in the size of replicates ( $m$  vs  $m+1$ ) and for overlapping units it would be more convenient to use a rate of overlap,  $k/m$ , so that there might not be exactly  $k$  units selected each time. We are in the process of doing simulation studies to evaluate the reduction in bias using this technique in more realistic conditions, and in the original problem of estimation of year-to-year change.

#### REFERENCES

- Cochran, W. (1977), *Sampling Techniques*, John Wiley & Sons, Inc., New York.
- Hinkins, S., Moriarity, C., and Scheuren, F. (1996), "Replicate Variance Estimation in Stratified Sampling with Permanent Random Numbers", *Proceedings Section on Survey Research Methods, American Statistical Association*.
- Wolter, K. (1985). *Introduction to Variance Estimation*, Springer-Verlag.